

This figure "fig1_ogle5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0006236v1>

This figure "fig2_ogle5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0006236v1>

This figure "fig3_ogle5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0006236v1>

This figure "fig4_ogle5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0006236v1>

This figure "fig5_ogle5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0006236v1>

This figure "fig6_ogle5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0006236v1>

This figure "fig7_ogle5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0006236v1>

This figure "fig8_ogle5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0006236v1>

Eclipsing Binaries in the OGLE Variable Star Catalogs. V. Long-Period β Lyrae-type Systems in the Small Magellanic Cloud and the PLC- β Relation

SLAVEK M. RUCINSKI

Electronic-mail: *rucinski@astro.utoronto.ca*

David Dunlap Observatory, University of Toronto
P.O.Box 360, Richmond Hill, Ontario, Canada L4C 4Y6

and

CARLA MACERONI

Electronic-mail: *maceroni@coma.mporzio.astro.it*

Rome Observatory, via Frascati 33, I-00040 Monteporzio C., Italy

ABSTRACT

Thirty eight long-period ($P > 10$ days), apparently contact binary stars discovered by the OGLE-II project in the SMC appear to be β Lyrae-type systems with ellipsoidal variations of the cool components dominating over eclipse effects in the systemic light variations and in the total luminosity. A new period–luminosity–color (PLC) relation has been established for these systems; we call it the PLC- β relation, to distinguish it from the Cepheid relation. Two versions of the PLC- β relation – based on the $(B - V)_0$ or $(V - I)_0$ color indices – have been calibrated for 33 systems with $(V - I)_0 > 0.25$ spanning the orbital period range of 11 to 181 days. The relations can provide maximum-light, absolute-magnitude estimates accurate to $\epsilon M_V \simeq 0.35$ mag. within the approximate range $-3 < M_V < +1$. In terms of their number in the SMC, the long-period β Lyrae-type binaries are about 50 times less common than the Cepheids. Nevertheless, their large luminosities coupled with continuous light variations make these binaries very easy to spot in nearby galaxies, so that the PLC- β relation can offer an auxiliary and entirely independent method of distance determination to nearby stellar systems rich in massive stars. The sample of the long-period β Lyrae systems in the SMC analyzed in this paper is currently the best defined and uniform known sequence of such binaries.

1. INTRODUCTION

This is the fifth paper based on the OGLE data, but the first one based on the data from the second phase of the OGLE project (OGLE-II, Udalski et al. 1997) utilizing the new 1.3 meter

Polish telescope on Las Campanas. The data are freely available over the computer networks¹.

Analysis of large numbers of eclipsing binary stars discovered by the OGLE-I project in the direction of Baade’s Window (BW) revealed that among almost one thousand binaries, about two thirds were contact systems of the W UMa-type (Rucinski 1997a, Rucinski 1997b = Papers I and II). The direction of the BW is towards the Galactic Bulge, yet all W UMa-type systems discovered there appear to be Disk population objects distributed over the long line of sight to the Bulge (Rucinski 1998b). While the W UMa-type binaries are very short-period systems, with orbital periods within $0.25 < P < 0.6$ days (this class of binaries is normally assumed to have periods shorter than one day), use of a Fourier decomposition on all available OGLE-I light curves showed that main-sequence contact binaries continue to exist in the Galactic Disk population to orbital periods of about 1.3 – 1.5 days and then abruptly vanish, with very few contact systems having longer periods (Rucinski 1998a = Paper III, see Figure 2 in this paper). Only four apparently contact systems were discovered with the orbital periods $P > 10$ days. Because of the long periods, these could not be main sequence stars such as those in the typical W UMa-type systems, but giants. Since there exist no theoretical predictions nor any observational indications that contact systems can be formed with giant components, no firm explanation for the long-period systems in BW could be offered at that time of Paper III. However, it was noticed that their light variations could be rather easily be explained by orbital synchronous rotation of a *single* distorted component filling its Roche lobe. Thus, these long-period binaries could consist of just one evolved component and some other star which – because of its small size and/or low temperature – would be invisible, yet would exert a sufficiently strong gravitational pull to distort the visible component.

In this paper we suggest that the “long-period contact binaries” are in fact representatives of the well known group of β Lyrae-type binaries. The primary basis for the argument is the well-populated sequence of long-period (LP) systems which has been established among contact binaries discovered by the OGLE-II project in the Small Magellanic Cloud (SMC)(Udalski et al. 1998a). We have found members of this sequence by using the same Fourier light-curve shape filter as in the previous papers of this series. Although the filter is meant to select genuine contact binary systems, it can also pass binaries which are not in contact, but which show sufficiently strong ellipsoidal variations and shallow or absent eclipses.

The hypothesis the LP systems are in fact binary systems with only one, Roche-lobe filling component dominating in the total light leads to a prediction of a period–luminosity–color (PLC) relation for such systems. Such a relation is expected to be very similar to what has been suggested for the W UMa-type stars (Rucinski 1994), but with only one component defining the global relationship. If the total mass and the mass-ratio are not correlated with the orbital period (however, this is an assumption only), a PLC relation will simply result from the Kepler-law scaling of the total systemic dimensions (and thus of the stellar radius) with the orbital period, $R \propto P^{2/3}$. Further conversion to a luminosity dependence, $L \propto R^2 T_{eff}^4$, and then to an absolute-magnitude

¹<http://www.astro.uw.edu.pl/~ogle/> or <http://www.astro.princeton.edu/~ogle/>.

dependence, will lead to $M_V \propto -10/3 \log P$. In this paper we show that a sample of β Lyrae systems in the SMC, selected with the exclusion of blue systems ($(V - I) > 0.25$), forms a surprisingly uniform and coherent group for which an absolute-magnitude relation – expressed as $M_V = M_V(\log P, color)$, with the expected $\log P$ dependence – can indeed be established.

We define the sample in Section 2 and then show the period–color relation (Section 3) as the best tool to delineate the sequence of the long-period contact systems. In Section 4 we show that in fact these are β Lyrae-type binaries, not genuine contact systems. The color-magnitude diagram (Section 5) leads directly to a period–luminosity–color which we express as a absolute-magnitude calibration (Section 6). We discuss the OGLE-II data for the systems discovered in the Baade’s Window sample in Section 7 and give estimates of the spatial frequency of the LP systems in Section 8. Section 9 summarizes the results of this paper.

2. SELECTION OF THE SAMPLE

The first step in the selection of the sample was finding the contact systems among all eclipsing binaries found by Udalski et al. 1998a in the SMC. Then, following the results of the next Section 3, which suggest a change in the sample characteristics at periods of about 10 days, the long-period representatives of the sequence were further selected. The $a_2 - a_4$ Fourier light-curve shape-filter has been applied to all eclipsing binaries listed in Udalski et al. 1998a. The operation of the filter is shown in Figure 1 while the details of the filter have been described before (Papers I, II, III and in Maceroni & Rucinski 1999 = Paper IV). The filter requires the $\cos(4\phi)$ term to be, in its absolute value, relatively small in relation to the $\cos(2\phi)$ term (ϕ is the orbital phase). A line on the $a_2 - a_4$ relation, pre-computed for a case of the inner contact using light curve synthesis models, defines a locus of systems with components just touching. Contact systems show small (in absolute value) a_4 coefficients when compared with a_2 . In effect, the Fourier light-curve filter selects eclipsing binaries with dominance of geometric distortion effects (which define the overall amplitudes and control the size of a_2) over eclipse effects (contributing primarily to the a_4 term).

Normally, the Fourier filter works very well in correctly identifying contact binaries. It also rejects very effectively semi-detached binaries of the Algol type which show deep eclipses. However, the filter may give wrong results for a semi-detached binary with very shallow or absent eclipses, in the situation of a great disparity of sizes of the components, with a large visible star showing a considerable tidal distortion². Then the a_4 coefficient can also be small relative to a_2 and the binary may be erroneously classified as a contact system. This is exactly what we suspect to be the case for the LP binaries described in this paper.

As an aside, we note that the OGLE-II project eliminated small amplitude systems from

²The filter fails also for systems with secondary minima displaced from phase 0.5. Such systems had to be removed manually from the sample.

consideration at the early stage by exclusion of the so-called “ellipsoidal variables” (Udalski et al. 1998a). This means that some small-amplitude contact, semi-detached and perhaps even detached binaries have been excluded at the stage of formation of the sample. We consider this choice as somewhat unfortunate because criteria for rejection were not clearly formulated in the discovery paper. Thus, the sample cannot be used for statistical studies involving light variation amplitudes or their derived quantities (such as a_2 and a_4) at the (poorly defined) low-amplitude ends of their distributions.

3. PERIOD – COLOR RELATION

The sequence of long-period contact systems is very well defined in the period-color (PC) relation which – as was many times stressed in this series – plays a special role for contact binary stars. The left panel of Figure 2 shows the PC relation for the all SMC systems selected as contact with the $a_2 - a_4$ light-curve shape filter. The sequence consists of 38 systems. It starts at 11 days (in fact a couple of systems with periods of 8 – 10 days may also belong to the sequence) and continues to 181 days. It has an inverted slope relative to the one normally observed for most common contact binaries of the W UMa-type: Typical contact binaries become hotter as their periods get longer, a trend which is explained by a progression upward along the main sequence. The inverted sense of the slope for the LP sequence suggests that they become cooler with the increase of dimensions, a property normally associated with giants.

The SMC sample is not the first one to show the LP sequence: In addition to the indications from the OGLE-I Baade’s Window (BW) sample that we mentioned in Section 1, the MACHO project has also detected several similar systems in LMC; the latter have been analyzed in Rucinski 1999. We show the LMC systems in the right panel of Figure 2. Although the BW and the LMC samples give a qualitative support to what will be discussed in this paper, we cannot utilize the BW and LMC data together with the SMC data for the following reasons: (1) It has been found that the OGLE-I photometry for BW suffers from zero-point offsets as well as from scale errors resulting from the non-linearity of the CCD detector. We discuss the details and present limited new OGLE-II data for three systems in Section 7; (2) The MACHO sample used unconventional spectral bandpasses which resulted in photometry only approximately transformable into the $(V, V - R)$ color system. Thus, neither the OGLE-I BW sample nor the MACHO sample can be compared with the new, high quality OGLE-II data for SMC.

4. LIGHT CURVES

The OGLE-II data for the SMC variable stars consist of light curves in the I band and of the maximum light magnitudes and color indices $(B - V)$ and $(V - I)$. The light curves are shown in Figures 3 and 4 while the photometric data are given in Table 1. The first columns of the

table contain the OGLE project data: First, the identification numbers (the consecutive ones used in this paper and then the OGLE-II numbers), followed by the orbital periods in days and the three photometric quantities, the I_{max} magnitude and the color indices $(B - V)$ and $(V - I)$. The OGLE-II identification number is made of the SMC field number (integer part of the number) and of the star number (six digits after the period). Among the 38 systems found in the SMC fields, three have additional identifications in the neighboring fields. The collection of the light curves in Figures 3 – 4 shows only the light-curve data based on the primary identifications. The light-curve Fourier decomposition coefficients a_i , which are given in the last six columns of Table 1, have been calculated for both identifications, but we used the primary ones in selection of the contact binary systems through the $a_2 - a_4$ Fourier filter. As in the previous papers of this series, a_i are the cosine coefficients while b_1 is the only sine term representing light curve asymmetries.

The light curves shown in Figures 3 and 4 are not of typical contact systems with very similar depths of eclipses. All show unequally deep eclipses which are characteristic for β Lyrae (EB) light curves. The light curves are well defined for most systems. Twenty seven systems with relatively large amplitudes of light variations have been selected to enter with full weight in determinations of luminosity calibrations in Section 6. Eleven systems with poorer light curves are marked by asterisks in Figures 3 – 4 and in Table 1; they are marked by open circles in the remaining figures.

The similarity of all LP systems in terms of their light curves is confirmed by the $a_2 - a_1$ plot in Figure 5 where we see a relatively tight relation, almost a proportionality between the coefficients. For normal W UMa-type systems, the a_1 coefficients are confined to a band $-0.02 < a_1 < 0$, without any clear correlation with a_2 (see Rucinski 1997b, Figure 5 there). Since the a_i coefficients are defined in terms of light intensity units, the proportionality of a_1 to a_2 implies similar light curves in magnitudes, with similar differences of the eclipse depths. This is exactly the property that was noticed when an attempt was made to model the light curve of the LP system in the BW sample, BW0.036 (Paper III): For a system where the only visible component fills the Roche lobe, the relative depths of eclipses are expected to be very similar almost irrespective of the orbital inclination and only very slightly depend on the mass-ratio.

Although the light curves are similar, they also show some small differences in that some show narrow secondary eclipses while some show relatively more scatter at both maxima. In one case, LP#9, we see well defined eclipses indicating that both components contribute to the systemic light; we show later that this is exactly the system which deviates from the PLC relation. We have made no attempt to model the light curves of the LP systems. We feel that without spectroscopic mass ratios, this would be futile exercise. We note that color curves cannot be currently formed because of much smaller numbers of the B and V observations relative to the number of I observations. However, all the light curve features can be explained by the same model which is applicable to the well-known prototype of β Lyrae.

At this point, we would like to make a clear distinction between the light-curve shape designated by EB from the name of close-binary stars β Lyrae. From now on, β Lyrae will be used

here as a binary-system prototype, not as a designator of the EB light-curve shape. That is, when calling a star a β Lyrae type, we will consider a binary with only one, giant-type component which dominates the total light, the other component being a main sequence star or collapsed star (perhaps surrounded by an accretion disk) with only a minor influence on the total systemic brightness and its light curve. Such a model is now well established for the prototype, following the extensive research of β Lyrae and of the related W Serpentis-type binaries (Huang 1963, Woolf 1965, Wilson 1974, Plavec 1985, Hubeny & Plavec 1991, Harmanec & Scholtz 1993, De Greve & Linnell 1994) and of various X-ray sources with giant-star mass donors. We stress that in calling a star a β Lyrae binary, we focus on two essential properties: the semi-detached model and the giant characteristics of the Roche-lobe filling component. Later on, to establish an absolute-magnitude calibration, we add the third property: the sufficiently yellow/red color which, in most cases, indicates a negligible photometric importance of the other component.

The light curves of the LP β Lyrae systems in the SMC sample appear to be symmetric, with identical heights of the light maxima (i.e. the so-called O’Connell effect is nonexistent for these systems). We also see no correlation between the sine coefficients, b_1 , and the coefficients measuring the eclipse-depth differences, a_1 . Such a correlation, manifesting itself as a preference for higher first maxima with the increased eclipse-depth difference, was observed for the short-period EB systems ($P < 1$ day), appearing in the Baade’s Window sample and analyzed in Rucinski 1997b. An explanation offered there was that these are semi-detached binaries with matter being transferred from the more massive, hotter to the less-massive, cooler components. Such systems are very different from the systems discussed in this paper in that they do not have to contain a giant, Roche-lobe filling component. In fact, judging by short orbital periods, many of them consist of main sequence stars first time evolving into the mass exchange phase or, perhaps, experiencing the semi-detached stage of the thermal-cycle relaxation oscillations (Lucy 1976, Flannery 1976, Robertson & Eggleton 1977, Lucy & Wilson 1979).

Finally, we note that the light curve amplitudes of the LP systems, as measured by the a_2 Fourier coefficient, do not show any correlation with the orbital period or the color index of the system. Note also that the light curve shapes in Figures 3 – 4 do not correlate in any obvious way with the periods (the periods are given in parentheses above each panel in these figures). In this respect, the sample appears to be uniform in the sense of consisting of very similar systems.

5. COLOR – MAGNITUDE DIAGRAM

The SMC data are an excellent material for forming a color-magnitude diagram (CMD) and for studying location of the LP systems in relation to the well recognized stages of stellar evolution. The distance modulus is relatively well known and will be soon known even better once the matter of systematic deviations between various methods of distance determinations to Magellanic Clouds is resolved (Udalski 1998, Udalski et al. 1998b, Gibson 1999, Nelson et al. 2000, Popowski 2000). Also, the average reddening for the SMC is small. In contrast, the Baade’s

Window sample (Paper III) suffered from large and uncertain reddening corrections and from the (then unrecognized) photometric errors, more fully described in Section 7. In addition, the distances of the BW systems were of course different precluding formation of an independent color-magnitude diagram (other than the one involving an absolute-magnitude calibration). In principle, the MACHO sample for the LMC (Rucinski 1999) could be used to form a CMD, but the data were in a non-standard photometric system and the reddening corrections (and their variations) are larger than typically for the SMC.

Figure 6 gives the observational CMD for the LP β Lyrae systems in the SMC, with the main stellar density enhancements (such as the main sequence or the red clump) marked by shading. The approximate discovery completeness limits of 90% (the slanting line starting at $I = 17.5$ and $V - I = 0$) and of close to 0% (1.5 mag. below), following Udalski et al. 1998a, are shown by slanting, dotted lines. In transforming from the observed, maximum-light photometric quantities to the de-reddened, absolute magnitudes, we used: $(m - M)_0 = 18.8$, $E_{V-I} = 0.12$ and $A_V = 2.5 E_{V-I}$. The LP systems are much more luminous than the main-sequence stars. They are also systematically brighter than typical giants in the SMC; this may be partly a selection effect as brightest stars are expected to have better photometry. Comparison with evolutionary tracks in Massey et al. 1995 indicates that the LP binaries considered in this paper cover the area normally associated with evolved, moderately massive ($3 - 5 M_\odot$) stars within the range of effective temperatures of $3.60 < \log T_{eff} < 4.00$. Later on, we set a blue limit at $(V - I)_0 = 0.25$ so that the effective temperature range is cut from above at $\log T_{eff} \simeq 3.9$.

The orbital periods of the LP β Lyrae systems are shown in Figure 6 by the size of the symbols. One immediately sees a period progression in that the systems with the longest periods appear in the upper part of the CMD. The longest periods, $P > 100$ days, appear exclusively among red giants above the Red Clump location, but a similar period progression is also visible in the upper-left, blue part of Figure 6 where five systems of similar color index have luminosities scaling with their orbital periods. This type of the period dependence indicates that a period – luminosity – color relation can be established for the LP systems.

6. PERIOD – LUMINOSITY – COLOR RELATION

6.1. Calibration of the relation

The previously established absolute magnitude calibrations for contact binary systems of the W UMa-type are of limited value here, primarily because of the vastly different ranges of the orbital periods. While the W UMa calibrations are applicable to the range 0.25 – 0.6 days (Rucinski & Duerbeck 1997)³, the calibration that we would like to establish will cover a large

³A blind application of the $(V - I)$ -based calibration of Rucinski & Duerbeck 1997 to the SMC LP systems gives M_V estimates about 3.5 magnitudes too bright.

range of the orbital periods from 11 days to 181 days.

We set aside and do not discuss the contact systems with the intermediate periods, starting around 0.6 – 1.5 days at the short end and extending to 10 days. Such systems apparently do not exist in the solar neighborhood and are exceedingly rare in the BW sample (Rucinski 1998a), but – possibly due to the discovery selection effect preferring brightest objects – are quite common in the present SMC and in the MACHO LMC samples (Figure 2). We feel that the intermediate-period (1.5 – 10 days) systems which are visible in the SMC and LMC samples are a mix of genuine contact systems with the moderately-short representatives of the β Lyrae systems discussed in this paper. They will be the subject of a separate investigation.

Assuming the same values for the distance modulus and reddening as in the previous section, the linear regression for the whole available data set of 38 systems (with 11 systems given half weights) gives:

$$M_V = -3.90 \log P + 3.27 (V - I)_0 + 2.40, \quad \sigma = 0.50 \quad (1)$$

$$M_V = -4.05 \log P + 3.01 (B - V)_0 + 3.46, \quad \sigma = 0.63 \quad (2)$$

The standard errors (per data point) σ are large and one may question the validity of linear fits in so poorly constrained cases. The deviations from the linear regressions are shown in the upper two panels of Figure 7. Note, that the bluest stars appear to deviate the most from the linear dependencies. If we exclude these systems setting the limit at $(V - I)_0 > 0.25$, then the linear fits improve substantially reaching the level of precision of about 0.35 mag. This level of accuracy is similar to that for W UMa-type binaries and for Cepheid variables. For the limited data set of 33 yellow-red systems, the linear weighted regression equations are:

$$M_V = -3.43 \log P + 2.04 (V - I)_0 + 2.80, \quad \sigma = 0.34 \quad (3)$$

$$M_V = -3.30 \log P + 1.49 (B - V)_0 + 3.38, \quad \sigma = 0.38 \quad (4)$$

The standard deviations are now quite respectably small so that the equations can be used for prediction of absolute magnitudes. The deviations from the linear fits for this case are shown in Figure 8. Realistic uncertainties of the coefficients for the sample of 33 yellow-red systems have been established by application of a bootstrap re-sampling process. The median values and the $1-\sigma$ ranges give the following relations:

$$M_V = -3.41^{+0.28}_{-0.27} \log P + 2.00^{+0.29}_{-0.31} (V - I)_0 + 2.80^{+0.27}_{-0.27} \quad (5)$$

$$M_V = -3.29^{+0.34}_{-0.34} \log P + 1.48^{+0.31}_{-0.34} (B - V)_0 + 3.36^{+0.43}_{-0.39} \quad (6)$$

The $\log P$ coefficients are very close what we suggested would be the case on the basis of very simplified considerations in Section 1, $M_V \propto -3.33 \log P$, which is highly gratifying. But we achieved that by removing the five blue systems with $(V - I)_0 < 0.25$. This removal is not entirely arbitrary; we discuss it in the next subsection. One can fix the period coefficient at the expected

Kepler-law value of -3.33 and determine the other two regression coefficients with higher much accuracy. The results for 33 yellow-red systems are:

$$M_V + 3.33 \log P = 1.94 (V - I)_0 + 2.72, \sigma = 0.34 \quad (7)$$

$$M_V + 3.33 \log P = 1.51 (B - V)_0 + 3.40, \sigma = 0.38 \quad (8)$$

while the bootstrap-estimated median values and the $1\text{-}\sigma$ ranges lead to:

$$M_V + 3.33 \log P = 1.94^{+0.14}_{-0.17} (V - I)_0 + 2.72^{+0.18}_{-0.14} \quad (9)$$

$$M_V + 3.33 \log P = 1.50^{+0.19}_{-0.20} (B - V)_0 + 3.41^{+0.16}_{-0.16} \quad (10)$$

We suggest that until better and more extensive data become available, Eqs. (3)–(4) or Eqs. (7)–(8) serve as the preliminary calibrations of the PLC- β relation.

6.2. Limitations of the PLC- β relation

One of the limitations of the PLC- β relation is the necessity to limit selection of the systems to redder than $(V - I)_0 = 0.25$. There are a few reasons why the blue stars may deviate from the general trends:

- The underpinning of the PLC- β relation is the assumption that the large, Roche-lobe filling component dominates the photometric characteristics of the system. Only then one can expect a tight relation between the size of this component (governed by the Roche lobe scaled with the size of the whole system) and the total luminosity of the binary system. Even a relatively small, but hot and hence possibly luminous companion can modify the color index to make it blue. Thus, blue color index may be considered as a direct indication that the hot companion of the Roche-filling star is contributing substantially to the system light. The well-observed system LP#9 (OGLE-II#3.134555) which shows clear eclipses is an excellent illustration of the point in that it is one of the blue systems removed from the final calibration.
- It is well known that the optical-range color indices, such as $(B - V)$ or $(V - I)$, lose sensitivity to effective temperature for very hot stars. The serious problems in the color – effective temperature transformations occur at $\log T_{eff} > 4.5$ (Massey et al. 1995). In our case, the blue stars which deviate from the PLC- β relation are not that hot, and have temperatures within the range $3.9 < \log T_{eff} < 4.0$.
- β Lyrae is itself a good case of a blue binary system which does not obey our PLC- β relation. In this case, it is not the contribution of the other component to the total light, but apparently the Roche-lobe filling component is too luminous. Dobias & Plavec 1985

summarize the parameters of relevance to our discussion: $P = 12.937$ days, $M_V = -4.7$ for both components and $M_V^{vis} = -4.1$ for the visible component, $(B - V)_0 = -0.13$ (which implies $(V - I)_0 \simeq -0.1$). Our calibrations predict $M_V^{VI} = -1.3$ and $M_V^{BV} = -0.5$ for the visible component. This component is thus some 2.8 – 3.6 mag. over-luminous relative to the calibrations.

- Possibly, use of single linear relations for the color dependence is simply inappropriate over the whole wide range $0 < (V - I)_0 < 1.6$ observed for the LP β Lyrae systems. The five hot systems in our sample do show the steep relation between M_V and $\log P$. If we lump all five and use their average color indices ($(V - I)_0 = +0.109$ and $(B - V)_0 = -0.067$) to enter these in Eqs. (3) and (4), then $M_V \propto -3.10 \log P$, with the zero points, $+0.87$ and $+1.19$ for the $(V - I)$ and $(B - V)$ based relations, respectively. These zero points are very different than for the more common yellow-red LP systems, leading to relations by about 2 to 2.5 magnitudes brighter. This would go some way towards explaining the over-luminosity of the β Lyrae less-massive component.

A serious limitation of any approach requiring de-reddened colour indices is the uncertainty of the reddening correction. Although the colour-index terms have relatively small coefficients and the strongest dependence is in the $\log P$ term, nevertheless the reddening corrections must be applied to the observed data. For the SMC OGLE-II data, two indices are available, $(B - V)$ and $(V - I)$. Can one use form a reddening-free index, similar to the well-known Q index? Unfortunately, this is not the case for the particular combination of the indices. An index constructed in the form: $Q' = (B - V) - k(V - I)$, where $k = E_{B-V}/E_{V-I}$ is not a monotonic function of the indices: It shows a shallow increase from $(V - I)_0 = -0.2$ to 0.9, then a broad maximum around $0.9 < (V - I)_0 < 1.4$, and a steep decline further to the red. Thus, while the conventional Q -index technique based on the $(U - B)$ and $(B - V)$ would work, the one based on the $(B - V)$ and $(V - I)$ indices would fail. The reddening correction uncertainty will remain the main limitation of the PLC- β relation.

6.3. External checks: The LMC and M31 data

Undoubtedly, calibrations of the PLC- β relation will be substantially improved when similar OGLE-II data for the Large Magellanic Cloud become available. At present, support from other sources is weak. As has been pointed before, the currently available MACHO data for the LP systems in LMC (Rucinski 1999) are in a non-standard photometric systems only approximating the VR filter system. Seven systems in this sample have orbital periods longer than 10 days, in the range $12 < P < 52$ days. Unfortunately, all seven are quite blue ($V - R < 0.35$, see their location in right panel of Figure 2). It is nevertheless possible to establish a tight linear relation, similar to the calibration Equations (3) – (10), for these systems. The relation ($M_V = -0.26 \log P + 4.31(V - R)_0 - 1.58$ for $m - M = 18.3$ and $E_{V-R} = 0.2$) has a small

dispersion $\sigma = 0.1$ mag., but we do not believe that this relation is of much validity because the photometric system is non-standard and all systems are blue.

The data for binary systems in the Andromeda galaxy detected by the DIRECT project also do not provide an external check on the calibration of the PLC- β relation. The project found four LP systems of β Lyrae-type in M31 (Stanek et al. 1998, Stanek et al. 1999, Kaluzny et al. 1999). All four have periods within 11 to 14 days, so that they do not probe the whole span of the orbital periods of the SMC sample. The DIRECT systems are: D31A-6423, D31C-13944, D31D-5186, D31D-7221 (we will call them below by consecutive numbers, 1 – 4). The systems #1 and #4 have very poor photometric data (especially the crucial color indices are poorly determined) while the system #2 has too blue color index (uncorrected $V - I = 0.13$) to belong to our proper LP sequence. Only the third system, D31D-5186 would qualify, with $P = 11.80$ days, $V_{max} = 19.5$, $V - I = 0.41$ and the very uncertain $B - V = -0.2$. The directly evaluated value of M_V , assuming $m - M = 24.5$ (Stanek & Garnavich 1998) and $E_{B-V} = 0.1$ is $M_V = -5.3$, while the one estimated from Eq. (3), assuming the same reddening, is $M_V = -0.2$. Thus, the discrepancy between the two estimates is very large, either shedding doubt on our SMC calibration or indicating that the companion is over-luminous (as in the very case of β Lyrae). We note that the DIRECT project can discover only the very brightest binaries in M31 and these would be the ones which would be most likely the ones which deviate the most from the calibration. Also, the color used is highly dependent on the assumed reddening which is only guessed in this case. We cannot exclude a possibility that $(V - I)_0$ for the system #3 is smaller than our threshold of 0.25; then the system would belong to the blue group which we explicitly excluded from consideration when establishing the PLC- β relation.

7. LONG-PERIOD SYSTEMS IN BAADE’S WINDOW

It has been realized recently that the OGLE-I photometry for Baade’s Window (BW) suffered from systematic zero point offsets and linearity problems resulting from the non-linearity of the original CCD detector. At the brightness level of the Bulge Red Clump Giants, the offsets are moderate: 0.021 in V , 0.035 in I and thus 0.056 in $(V - I)$ (Paczynski et al. 1999). More worrisome are however the non-linearities, relatively larger in the V -filter data for red stars and affecting particularly strongly the $(V - I)$ color indices of faint and red stars. Dr. Udalski (private communication) estimated that deviations in $(V - I)$ may be as large as 0.2 mag. for faintest surveyed stars. Since the color indices play an important role in our absolute-magnitude calibrations, the deficiency of the OGLE-I photometry will require a more extensive re-discussion of the results of Papers II and III.

The OGLE project, currently in its Phase II, is characterizing the systematic errors of OGLE-I. Some stars already have excellent new data in V , $(B - V)$ and $(V - I)$. At our request, Dr. Udalski kindly checked the availability of new data for the sequence of the systems with $P > 1$ days which had been discussed in Rucinski 1998a (Paper III). About two thirds of these

stars have new photometry. Among the four BW systems with $P > 10$ days, three have new photometric data. These have been made available to us in terms of the average light levels, requiring additional corrections to the maximum light level. They are based on typically 30 observations in I and 10 observations in V . We applied the differences $\Delta I = \overline{I_1} - I_1$, as found on the basis of the OGLE-I light curves (Paper III), to convert the data from the average ($\overline{I_2}$) to the maximum (I_2) light levels. The details are given in Table 2.

The left panel of Figure 2 shows the OGLE-II data for the three LP β Lyrae systems in the period-color plane, with the assumed reddening corrections the same as in Paper III (the three systems are marked by filled asterisks). The corrections were found on the basis of an assumption that the reddened stars are further than 2 kpc and thus suffer the same amount of reddening as the Bulge stars in the BW (Stanek 1996). This assumption can be checked for consistency using the calibration in Eq. (3). The estimates of the distances are given in the last column of Table 2. The errors reflect the uncertainty of the calibration equation of about 0.3 mag. The results are consistent with two systems in the Bulge (possibly on the other side of it) and one system possibly in front of the Bulge or inside its elongated part directed toward us. The fourth system, #0.036, without the new data, also appears to be located in the Bulge. We treat these distance determinations as very preliminary and disregard a possible metallicity mismatch between the calibration established for the SMC systems and the possibly metal-rich systems in Baade’s Window.

8. SPATIAL FREQUENCY

The spatial frequency of the long-period β Lyrae-type systems is difficult to estimate because it is practically impossible to estimate the OGLE-II search volume in the SMC. In terms of the intrinsic luminosities, the LP β Lyrae-type systems occupy the range $-5 < M_V < +1$ (the calibrated part extends to $M_V \simeq -3$) so that they can be compared with the Cepheid variables. The OGLE-II project observed 2155 Cepheids in the SMC (Dr. A. Udalski – private information) which can be compared with 38 long-period β Lyrae-type systems discussed here. While a large fraction were previous discoveries and the LP binary systems are new, this gives a rough idea about the frequency of occurrence of the LP β Lyrae-type more than 50 times below the spatial frequency of the Cepheids. As an aside, notice when comparing the two types of variables, that the period term in the LPC- β relation appears to be close to the expected $M_V \propto -3.33 \log P$, while the Cepheids show a period dependence, with the period coefficient -2.78 (Feast & Walker 1987) or -2.76 (Madore & Friedman 1991). The similarity of these dependencies should not be surprising as both are driven by the same dimension – response time relationship of the dynamical time scale.

The Baade’s Window data give us only a very approximate constraint on their frequency in our Galaxy. If one determines the average frequency of occurrence by simply dividing their number (four) by the volume of the OGLE-I search in the BW area (about 2.5×10^7 pc³ to the

Galactic center), then an estimate of the frequency is 1.6×10^{-7} LP systems per cubic parsec. However, the LP β Lyrae systems appear to be strongly concentrated towards the Galactic Bulge so that such a naive estimate does not take into account the increase in numbers of stars as the line of sight penetrates the Galactic Bulge. We recall that the W UMa-type systems of the disk population approximately follow the stellar density in the BW direction with the average density of about 7.6×10^{-5} systems per cubic parsec (Rucinski 1998b).

9. DISCUSSION

This paper discusses consequence and observational support for an assumption that the long-period ($P > 10$ days), apparently contact binary systems observed in Baade’s Window, and in LMC and SMC, are in fact β Lyrae systems with the cool components dominating their total luminosity. We analyzed in detail the LP systems in the SMC and found that they form a well-defined, moderately tight, “inverted” period-color relation, implying a color–size relation characteristic for giants rather than for main-sequence stars. The principal support, however, comes from the existence of a period-color-luminosity relation for those which are sufficiently cool. Tentatively, we set the color-index limit at $(V - I)_0 > 0.25$. An absolute-magnitude calibration of the new PLC- β relation in the form $M_V = M_V(\log P, color)$, established in a preliminary form on the basis of 33 systems in the SMC, has the expected slope $M_V \propto -3.33 \log P$. It offers an exciting application for distance determination to nearby galaxies. The LP β Lyrae systems are the brightest among binary stars and their continuous light variations make them also very easy to spot. They occupy roughly the same range of the absolute magnitudes as the Cepheid variables, but appear to be by about 50 times less common in space. Therefore, the PLC- β relation can play only an auxiliary – although entirely independent – role in distance determinations.

We see sequences of ostensibly contact, long-period systems not only in the SMC, but also in Baade’s Window and in MACHO data for LMC. However, the latter data sets could not be used to improve the absolute magnitude calibration of the PLC- β relation for reasons described in detail in Sections 3 and 7. It is expected that the data for LMC currently collected by the OGLE-II project will provide an excellent check and improvement of the tentative results presented here.

We conclude by observing that the β Lyrae binaries discussed in this paper are expected to show some similarities to the cataclysmic variable (CV) binaries. The CV binaries are also known to have secondary (less massive components) filling their Roche lobes. As was observed by Faulkner et al. 1972, a combination of Kepler’s law, $a^3 \propto (M_1 + M_2) P^2$, with the approximate relation for the relative radius of the Roche lobe filling star, $R_2/a = 0.462(q/(1 + q))^{1/3}$, with $q = M_2/M_1 \leq 1$, gives a relation similar to that for pulsating stars: $P = 0.438/\sqrt{\rho}$, where the period P is in days and the mean density of the star, ρ , is in g cm^{-3} . This relation is valid for small mass ratios ($q < 0.5$), but corrections for larger values of q are small. Because the range of the observed period for the LP sequence in the SMC is large, one expects the Roche-lobe filling components to have mean densities within a large range from about $2 \times 10^{-3} \text{ g cm}^{-3}$ to less than

$10^{-5} \text{ g cm}^{-3}$, that is within typical range observed for yellow-red giants and supergiants.

Special thanks are due to Dr. Andrzej Udalski for his prompt response – in spite of many ongoing activities – to a request to extract new OGLE-II data for the LP systems observed by OGLE-I project in Baade’s Window and for several explanatory comments on the data. Thanks are also due to Dr. Stefan Mochnacki for very useful comments and suggestions. The authors would like to thank the OGLE team for making the data freely available over the computer networks.

CM acknowledges the support of the Italian Ministry of Research (MURST) through a Cofin98 grant and SMR acknowledges the support of the Natural Sciences and Engineering Research Council (NSERC) of Canada.

REFERENCES

- De Greve, J.P., & Linnell, A.P. 1994, *A&A*, 291, 786
- Dobias, J.J. & Plavec, M.J. 1985, *AJ*, 90, 773
- Faulkner, J., Flannery, B.P., & Warner, B. 1972, *ApJ*, 175, L79
- Feast, M.W. & Walker, A.R. 1987, *ARA&A*, 25, 345
- Flannery B.P., 1976, *ApJ*, 205, 217
- Gibson, B.K. 1999, *Mem.Soc.Astr.Ital.*, in press; astro-ph/9910574
- Harmanec, P., & Scholtz, G. 1993, *A&A*, 279, 131
- Hunag, S.–S. 1963, *ApJ*, 138, 342
- Hubeny, I., & Plavec, M.J. 1991, *AJ*, 102, 1156
- Kaluzny, J., Mochejska, B.J., Stanek, K.Z., Krockenberger, M., Sasselov, D.D. Tonry, J.L., & Mateo, M. 1999, *ApJ*, 118, 346
- Lucy L. B., 1976, *ApJ*, 205, 208
- Lucy L. B., Wilson R. E., 1979, *ApJ*, 231, 502
- Maceroni, C., & Rucinski, S.M. 1999, *AJ*, 118, 1819 (Paper IV)
- Madore, B.F. & Friedman, W.L. 1991, *PASP*, 103, 667
- Massey, P., Lang, C.C., DeGioia-Eastwood, K., & Garmany, C.D. 1995, *ApJ*, 438, 188

- Nelson, C.A., Cook, K.H., Popowski, P., & Alves, D.R. 2000, *AJ*, 119, 1205
- Paczynski, B., Udalski, A., Szymanski, M., Kubiak, M., Pietrzynski, G., Soszynski, I., Wozniak, & P. Zebrun, K. 1999, *Acta Astron.*, 49, 319
- Plavec, M.J. 1985, in *Interacting Binaries*, edited by P.P. Eggleton and J.E. Pringle (Reidel, Dordrecht), p. 155
- Popowski, P. 2000, *ApJ*, 528, L9
- Robertson J.A., Eggleton P.P., 1976, *MNRAS*, 179, 359
- Rucinski, S.M. 1994, *PASP*, 106, 462
- Rucinski, S.M. 1995, *PASP*, 107, 648
- Rucinski, S.M. 1997a, *AJ*, 113, 407 (Paper I)
- Rucinski, S.M. 1997b, *AJ*, 113, 1112 (Paper II)
- Rucinski, S.M. 1998a, *AJ*, 115, 1135 (Paper III)
- Rucinski, S.M. 1998b, *AJ*, 116, 2998
- Rucinski, S.M. 1999, *Acta Astr.*, 49, 341
- Rucinski, S.M., & Duerbeck, H.W. 1997, *PASP*, 109, 1340
- Stanek, K.Z. 1996, *ApJ*, 460, L37
- Stanek, K.Z., & Garnavich, P.M. 1998, *ApJ*, 503, L131
- Stanek, K.Z., Kaluzny, J., Krockenberger, M., Sasselov, D.D. Tonry, J.L., & Mateo, M. 1998, *ApJ*, 115, 1894
- Stanek, K.Z., Kaluzny, J., Krockenberger, M., Sasselov, D.D. Tonry, J.L., & Mateo, M. 1999, *ApJ*, 117, 2810
- Udalski, A. 1998, *Acta Astron.*, 48, 113
- Udalski, A., Kubiak, M., & Szymanski, M. 1997, *Acta Astron.*, 47, 319
- Udalski, A., Soszynski, I., Szymanski, M., Kubiak, M., Pietrzynski, G., Wozniak, P., & Zebrun, K. 1998a, *Acta Astron.*, 48, 563
- Udalski, A., Pietrzynski, G., Wozniak, P., Szymanski, M., Kubiak, M., & Zebrun, K. 1998b, *ApJ*, 509, L25
- Wilson, R.E. 1974, *ApJ*, 189, 319

Woolf, N.J. 1965, ApJ, 141, 155

Captions to figures:

Fig. 1.— The Fourier light-curve shape filter selects contact binaries by comparison of the a_2 and a_4 cosine coefficients. Systems with properties of contact binaries are located below the curved line. The small dots give the data for all eclipsing systems in the SMC while the large filled circles mark 38 long-period ($P > 10$ days) contact systems discussed in this paper. The open circles are used for 11 systems with relatively poorer light curves which have been given half weight in luminosity calibrations discussed further in the paper. Note that contrary to the results for Baade’s Window (Rucinski 1997a), the contact domain is rather thinly populated in the SMC, mostly due to the absence of the genuine W UMa-type binaries ($P < 1$ day). We give arguments in the paper that the apparently contact long-period systems are in fact semi-detached binaries of the β Lyrae-type with component-distortion effects dominating over shallow or absent eclipses.

Fig. 2.— The period–color diagram for all contact systems in the SMC (this paper, left panel) and LMC (the MACHO data, right panel, as discussed in Rucinski 1999). The large circles show systems discussed in this paper; the open circles mark those which have been assigned half weights in luminosity calibrations discussed later on. The SMC data are based on the $(V - I)$ color index while the LMC data are based on the $(V - R)$ color index. The curves give the blue short-period envelope for normal, short-period W UMa-type systems. For simplicity, we have assumed the same reddening vectors for all systems in the SMC, $E_{V-I} = 0.12$ (vertical lines). In subsequent sections we find that systems with $(V - I)_0 > 0.25$ provide a well defined period – luminosity – color relation. Three large asterisks show locations of the long-period systems in Baade’s Window; they are more fully discussed in Section 7. We cannot directly utilize the BW or LMC data in this paper, but point out the existence of the LP systems which appear to be in contact and form a sequence extending to long orbital periods.

Fig. 3.— A collection of I band light curves for 38 long period contact systems in the SMC. This figure contains 24 light curves; the remaining curves are in the next figure. The systems are identified by numbers above each of the panels with the orbital periods given in parentheses. An asterisk added to the star number indicates half-weight given in the luminosity calibrations. The vertical scale of each panel spans one magnitude and extends from 0.6 mag. below to 0.4 mag. above the average magnitude. The maximum light magnitudes, the orbital periods, the color data as well as the Fourier cosine coefficients are given in Table 1.

Fig. 4.— The same as for Figure 3, but for 14 among 38 long period contact systems in the SMC.

Fig. 5.— The first cosine term of the light-curve Fourier decomposition, a_1 , appears to scale in proportion to the largest term, a_2 . This confirms that the light curves are very similar in shape. The open circles mark systems with half weights in the calibration solutions.

Fig. 6.— The color-magnitude diagram for the LP systems in SMC. The orbital periods of individual systems are marked by the size of the circle. The right and upper axes show the dereddened color index and absolute magnitude data for $E_{V-I} = 0.12$ and $(m - M)_0 = 18.8$. The

slanted dotted lines give the approximate locations of the 90% and 0% completeness levels of the OGLE-II variable star search. The broken lines represent the constant period lines according to the calibration of our PLC- β relation as in Eq. (7). They extend to the red of $(V - I)_0 = 0.25$. The current calibrations are not valid to the blue of that point (marked by the vertical dotted line), although a period dependence with a correct slope is observed for such systems as well. The main sequence (MS) and the red clump (RC) of the SMC are marked by shading.

Fig. 7.— The deviations from the regression lines of the calibration in Equations (1) – (2). Note the large negative deviations at blue color indices. The labels are the consecutive numbers in Table 1; they are given for systems deviating more than 0.75 mag. from the linear regression. Note in particular the location of the system LP#9 which is discussed in the text. The dotted lines give the color-index cutoff lines at $(V - I)_0 = 0.25$ and the corresponding $(B - V)_0 = 0.21$.

Fig. 8.— Same as in Figure 7, but with a limitation to systems with $(V - I)_0 > 0.25$, as in Equations (3) – (4). Notice a small inconsistency – which we disregarded in our determinations of the PLC calibrations – in that 3 systems are slightly bluer than $(B - V)_0 = 0.21$ which roughly corresponds to the cutoff at $(V - I)_0 = 0.25$.

TABLE 1. β Lyrae systems with $P > 10$ days in the Small Magellanic Cloud

LP#	SMC#	P(d)	I_{max}	$(B - V)$	$(V - I)$	a_0	a_1	a_2	a_3	a_4	b_1
1	1.012977	60.1884	14.279	0.020	0.257	0.859	-0.043	-0.129	-0.024	-0.011	-0.005
2	1.035641	29.1173	17.711	0.378	0.735	0.936	-0.015	-0.068	-0.002	0.002	-0.002
3*	1.062436	13.6637	18.351	0.768	1.039	0.972	-0.022	-0.041	0.003	0.004	-0.007
4	3.002900	89.8238	16.183	1.251	1.372	0.939	-0.011	-0.064	-0.011	0.002	-0.006
5	3.060633	28.2032	17.894	0.752	0.988	0.873	-0.025	-0.114	-0.015	-0.022	-0.004
6*	3.075910	72.9871	16.928	1.003	1.177	0.956	-0.032	-0.029	0.000	0.000	-0.003
7	3.116114	24.3557	17.953	0.386	0.795	0.921	-0.019	-0.089	-0.017	-0.008	-0.002
8*	3.116383	13.2463	19.070	0.216	0.827	0.968	0.018	-0.082	0.006	-0.011	-0.003
9	3.134555	11.8497	16.283	0.129	0.333	0.845	-0.072	-0.173	-0.028	-0.039	-0.005
10*	3.134806	55.5766	17.898	...	1.310	0.899	-0.019	-0.099	-0.011	-0.011	-0.001
11*	3.149100	33.9911	18.028	0.750	1.023	0.937	-0.003	-0.050	0.000	-0.002	0.008
12	3.193411	28.4187	17.197	0.195	0.555	0.917	-0.028	-0.112	-0.016	-0.026	0.003
13	3.238378	43.8842	14.880	-0.047	0.158	0.957	-0.009	-0.053	-0.008	-0.001	-0.005
14	4.156495	22.2481	17.174	0.219	0.435	0.899	-0.025	-0.123	-0.011	-0.023	0.001
15*	4.179449	20.3446	18.088	0.987	1.275	0.951	-0.020	-0.055	-0.006	-0.005	-0.005
16	5.145588	138.5914	15.730	1.056	1.349	0.881	-0.044	-0.122	-0.028	-0.025	0.000
17	5.162500	181.0044	15.729	1.583	1.638	0.928	-0.021	-0.083	-0.022	-0.010	-0.002
18	5.180039	97.1930	16.065	0.769	1.041	0.907	-0.017	-0.074	-0.006	-0.007	0.000
19*	5.257236	11.7916	18.788	0.347	0.778	0.915	-0.016	-0.110	-0.018	-0.019	-0.009
	(6.006812)	11.7957	18.790	0.346	0.755	0.934	-0.027	-0.074	-0.014	-0.005	-0.010
20	6.018257	15.5577	18.609	0.536	0.805	0.883	-0.035	-0.150	-0.023	-0.032	-0.003
21	6.094503	112.4580	15.290	0.799	0.922	0.893	-0.040	-0.113	-0.031	-0.021	0.009
22	6.110581	88.3550	16.046	1.067	1.171	0.871	-0.056	-0.162	-0.036	-0.027	-0.005
23	6.122285	28.5879	15.899	0.008	0.117	0.905	-0.012	-0.095	-0.009	-0.005	0.000
24*	6.199624	62.1454	16.547	1.000	1.067	0.972	-0.019	-0.030	0.001	-0.001	0.002
25	6.232915	13.1408	18.136	0.338	0.422	0.897	-0.032	-0.118	-0.014	-0.017	0.006
26	6.292000	21.8606	18.049	0.916	1.147	0.937	-0.049	-0.104	-0.003	-0.020	-0.019
27	6.297249	21.1419	18.082	0.406	0.850	0.923	-0.023	-0.089	-0.014	-0.004	-0.005
28	7.018931	12.6850	18.497	0.352	0.599	0.905	-0.023	-0.110	-0.023	-0.019	-0.001
29	7.255665	47.3855	16.626	0.813	1.066	0.875	-0.037	-0.156	-0.027	-0.035	-0.003
	(8.030691)	47.4408	16.628	...	1.053	0.861	-0.026	-0.152	-0.026	-0.038	-0.008
30	7.263170	35.6454	17.343	0.576	0.902	0.858	-0.042	-0.156	-0.021	-0.040	0.004
	(8.038913)	35.6472	17.352	0.583	0.884	0.865	-0.032	-0.162	-0.011	-0.048	0.006
31*	8.000105	31.8424	17.068	0.817	1.075	0.970	-0.012	-0.042	-0.004	-0.004	-0.012
32	8.015451	29.4313	18.226	0.737	0.796	0.873	-0.033	-0.152	-0.016	-0.042	0.003
33	9.059110	16.3410	17.739	0.657	0.905	0.924	-0.012	-0.094	-0.010	-0.012	-0.005
34*	9.061975	25.5624	18.100	0.760	1.066	0.981	-0.037	-0.042	0.002	-0.007	-0.004
35	10.059436	52.8010	17.313	0.946	1.182	0.906	-0.010	-0.116	-0.004	-0.026	-0.002
36	10.078655	129.0291	15.308	0.980	1.156	0.881	-0.050	-0.141	-0.029	-0.031	-0.002
37	10.088122	14.3973	16.444	0.057	0.282	0.895	-0.017	-0.094	-0.019	-0.019	-0.009
38*	11.005338	12.0634	18.431	0.309	0.577	0.943	-0.004	-0.098	-0.003	-0.017	-0.011

Notes to Table 1.

The SMC OGLE-II identifications are given in the first column as the field number (integer part) and star number (fractional part). In three cases identifications in the neighboring fields were available; these are given in parentheses following the primary identification. The $a_2 - a_4$ data of the primary identifications have been used as criteria for inclusion in this table. Asterisks in the first column identify systems which have been given half weight in the calibrations of the PLC- β relation. The de-reddened, absolute-magnitude data used in the paper have been derived using: $V_{max} = I_{max} + (V - I)$, $(V - I)_0 = (V - I) - E_{V-I}$, $M_V = V_{max} - (m - M)_0 - A_V$, where $(m - M)_0 = 18.8$ and $A_V = 2.5E_{V-I}$, with $E_{V-I} = 0.12$. For more detail, consult the text.

TABLE 2. OGLE-II Data for β Lyrae systems with $P > 10$ days in Baade's Window

OGLE-I	OGLE-II	P(d)	E_{V-I}	$\bar{I}_2$	$\overline{(V-I)}_2$	ΔI_1	I_1	$(V-I)_1$	I_2	V_2	M_V^{cal}	d(kpc)
0.036	...	26.307	0.70	...	...	...	15.51	1.82	...	(17.33)	(0.21)	(11.8 ± 1.8)
3.051	45.501074	14.243	0.77	16.08	1.88	0.05	15.98	1.83	16.03	17.91	1.11	9.4 ± 1.2
3.079	46.036352	12.313	0.87	16.51	2.06	0.05	16.43	2.07	16.46	18.52	1.49	9.3 ± 1.2
4.002	46.135504	22.672	0.67	14.59	1.79	0.18	14.33	1.99	14.41	16.20	0.44	6.6 ± 0.8

Notes to Table 2.

The first two columns give the star identifications consisting of the respective OGLE field numbers (integer part) and of the number in the field (fractional part). The reddening estimate for $d > 2kpc$, E_{V-I} , and the maximum light values of I_1 and $(V-I)_1$ are from Paper III. The differences between the mean and maximum levels, ΔI_1 , have been calculated on the basis of the OGLE-I light curves. The values of ΔI_1 have been used to convert the average OGLE-II data to maximum light level values, I_2 . It has been assumed that the color indices $(V-I)_2$ are the same for the average and maximum light levels. The values of M_V^{cal} have been calculated using Eq. (3) leading to the distance estimates through: $V - M_V^{cal} = 5 \log d - 5 + 2.5 E_{V-I}$. The OGLE-II data are not available for the system #0.036; in this case the distance estimate is based on the OGLE-I data.